
No.



200300207

THE UNITED SHATES OF ANTERIOS

TO ALL TO WHOM THESE: PRESENTS SHALL COME:

Ilorida Agricultural Experiment Station

LOCCUS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE $\mathbf{L}\mathbf{A}\mathbf{W}$.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR ING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE URPOSE, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT THE PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY OLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMB ERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321

PEANUT

'Hull'

In Testimonn Thereof, I have hereunto set my hand and caused the seal of the Hunt Unriety Protection Office to be affixed at the City of Washington, D.C. this twenty-fifth day of August, in the year two thousand and five.

ET SEO

Plant Varioly Protection Off

REPRODUCE LOCALLY. Include form number and date on all	reproductions.	·	FORM APPROVED - OMB NO. 0581-0055
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTEC	The following statements are ma 1974 (5 U.S.C. 552a) and the Pa	ade in accordance with the Frivacy Act of aperwork Reduction Act (PRA) of 1995.	
APPLICATION FOR PLANT VARIETY PROTECTION Unstructions and information collection burden statements	Application is required in order to determine if a plant variety protect certificate is to be issued (7 U.S.C. 2421). Information is held confide until certificate is issued (7 U.S.C. 2426).		
NAME OF APPLICANT(S) (as it is to appear on the Certificate)	nt on reverser	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
Florida Agricultural Experiment Stati	.on	UF98326	Hull
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Count	η)	6. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY
Office of Dean for Research 1022 McCarty Hall, University of Flor		352-392-1784	2 0 0 3 0 0 2 0 7
P. O. Box 110200 Gainesville, FL 32611-0200		6. FAX (include area code)	F DATE
Gainesville, II J2011-0200		352-392-4965	March 31,2003
7. GENUS AND SPECIES NAME	·		N G
	8, FAMILY NAME (Bota		FILING AND EXAMINATION FEE:
Arachis hypogaea L.	Legumino	saea	F 3652 -
9. CROP KIND NAME (Common name)			1 March 31,2003
Peanut (Groundnut)			E
10. If THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZAT	ION (corporation, partners	hip, association, etc.) (Common name)	E 432
Florida Agricultural Experiment Stati 11. IF INCORPORATED, GIVE STATE OF INCORPORATION	.011		
		12. DATE OF INCORPORATION	Date June 28, 2005
NA 13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERV		NA ·	14. TELEPHONE finclude area codel
Dr. D. W. Gorbet North Florida Research and Education 3925 Highway 71 Marianna, FL 32446	granical section		15. FAX (include area code)
16. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow in.	etropiano en Escapa.		
a. Exhibit A. Origin and Breeding History of the Variety	Su ocuons on reverse,		
b. XI Exhibit B. Statement of Distinctness			
c. Exhibit C. Objective Description of the Variety			•
d. X Exhibit D. Additional Description of the Variety (Optional)			
e. 🖾 Exhibit E. Statement of the Basis of the Applicant's Ownership f. 🖾 Voucher Sample (2,600 viable untreated seeds or, for tuber propagated		al	in dia agrand gublic repositore
g. Siling and Examination Fee (\$2,460), made payable to "Tressurer of the			mea m ап арргочев равно герозкогу;
17. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY V XI YES (If "yes," answer items 18 and 19 below)		A CLASS OF CERTIFIED SEED? (See See	tion 83(a) of the Plant Variety Protection Acti
18. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LIMITED A GENERATIONS?	S TO NUMBER OF	9. IF "YES" TO ITEM 18, WHICH CLASSI	ES OF PRODUCTION BEYOND BREEDER SEED?
M YES □ NO		☑ FOUNDATION ☑ REGISTE	RED KI CERTIFIED
20. HAS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY BEEN RELE	ASED, USED, OFFERED F	OR SALE, OR MARKETED IN THE U.S. OF	R OTHER COUNTRIES?
USA, May 2002			
21. The applicant(s) declare that a viable sample of basic seed of the variety will be applicable, or for a tuber propagated variety a tissue culture will be deposited in	furnished with application a public repository and m	and will be replenished upon request in ac aintained for the duration of the certificate	cordance with such regulations as may be
The undersigned applicant(s) is(are) the owner(s) of this sexually reproduced or t Section 42, and is entitled to protection under the provisions of Section 42 of the	luber propagated plant var Plant Variety Protection	icty, and believe(s) that the variety is new Act.	, distinct, uniform, and stable as required in
Applicant(s) is(are) informed that false representation herein can jeopardize prote	ction and result in penalti	23.	
SIGNATURE OF APPUCANT (Owner(s))	SIGNAT	URE OF APPLICANT (Orghor(s))	
NAME (Please print or type)	NAMET	print or type)	
D. W. Gorbet	Ri	chard L. Jones	
CAPACITY OR TITLE DATE	}	TY OR TITLE	DATE
Professor/Breeder	De	ean for Research	
STD-470 (03-96) (Previous editions are to be destroyed)	and the second section of the second	(See reverse for instructions an	d information collection burden statement)

REPRODUCE LOCALLY, Include form number and date on all I	reproductions		Form Approved - OMB No. 0581-0055
U.S. DEPARTMENT OF AG AGRICULTURAL MARKETII SCIENCE AND TECHNOLOGY - PLANT VAF	MC CEDVICE	The following statements are made in acc the Paperwork Reduction Act (PRA) of 1	ordance with the Privacy Act of 1974 (5 U.S.C. 552a) an 195.
APPLICATION FOR PLANT VARIETY PR (Instructions and information collection but	OTECTION CERTIFICATE	Application is required in order to determing (7 U.S.C. 2421). Information is held confi	ne if a piant variety protection certificate is to be Issued dential until certificate is issued (7 U.S.C. 2426).
1. NAME OF OWNER Florida Agricultural Experi	ment Station RAW	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NAME	3. VARIETY NAME
University of Florida, IFAS	5/6/01		Hull
4. ADDRESS (Street and No., or R.F.D. No., City, State, and	ZIP Code, and Country)	5. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY
Office of Dean for Research 1022 McCarty Hall, Universi P. O. Box 110200	ty of Florida	352-392-1784 6. FAX (include area code)	PVPO NUMBER
Gainesville, FL 32611-0200	and the second	352-392-4965	FUNCELEE
 IF THE OWNER NAMED IS NOT A "PERSON", GIVE FOR ORGANIZATION (corporation, partnership, association, etc 	M OF 8. IF INCORPORATED, GIVE STATE OF INCORPORATIO	9. DATE OF INCORPORATION	FILING DATE
University Ag. Expt. Stn. (Pul	olic) NA	NA	
10. NAME AND ADDRESS OF OWNER REPRESENTATIVE	S) TO SERVE IN THIS APPLICATION. (Fi	rst person listed will receive all papers)	F FILING AND EXAMINATION FEES:
Dr. Daniel W. Gorbet North Florida Research and 3925 Highway 71 Marianna, FL 32446	Education Center		E S S DATE E CERTIFICATION FEE:
			V E DATE
11. TELEPHONE (Include area code)	12. FAX (Include area code)	13. E-MAIL	14. CROP KIND (Common Name)
850-482-9956	850-482-9917	dgorbet@mail.ifas.ufl.edu	Peanut
15. GENUS AND SPECIES NAME OF CROP		16. FAMILY NAME (Botanical)	17. IS THE VARIETY A FIRST GENERATION
Arachis hypogaea L.		Leguminoseae	HYBRID?
18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT (Follow instructions on reverse) a. 区 Exhibit A. Origin and Breeding History of the Various Exhibit B. Statement of Distinctness	· · · · · · · · · · · · · · · · · · ·	19. DOES THE OWNER SPECIFY THAT SET CERTIFIED SEED? See Section 83(a) YES (If "yes", answer items 20 a	
c. 🛛 Exhibit C. Objective Description of Variety		VARIETY BE LIMITED AS TO NUMBER O	OF CLASSES?
d. X Exhibit D. Additional Description of the Variety (O	ofional)	IF YES, WHICH CLASSES? 🗓 FOU	NDATION A REGISTERED K CERTIFIED
e.	for tuber propagated varieties, I maintained in an approved public	21. DOES THE OWNER SPECIFY THAT SEE VARIETY BE LIMITED AS TO NUMBER OF FYES, SPECIFY THE NUMBER 1,2,3, et	D OF THIS IN YES IN NO DE GENERATIONS?
22. HAS THE VARIETY (INCLUDING ANY HARVESTED MATE FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRAN OTHER COUNTRIES? A YES May 2002 IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (FIRST)	SFERRED, OR USED IN THE U. S. OR NO EDISCOSITION TRANSFER OF 195	23. IS THE VARIETY OR ANY COMPONENT PROPERTY RIGHT (PLANT BREEDER'S USA-Utility Patents A Yes	OF THE VARIETY PROTECTED BY INTELLECTUAL RIGHT OR PATENT)? on oil chemistry In No OF FILING OR ISSUIANCE AND ASSIGNED
24. The owners declare that a viable sample of basic seed of for a tuber propagated variety a tissue culture will be deposited to a tuber propagated owner(s) Is(are) the owner of this sexual and is entitled to protection under the provisions of Section	lly reproduced or tuber proposed at all at	to for the duration of the certificate.	language of the Salata Araba and the Company of the
Owner(s) is(are) informed that false representation herein		nallies.	
SIGNATURE OF DIVINER	1	SIGNATURE OF OWNER	<u> </u>
11 _M		SIGNATURE OF CLANER	
- Land WA	rlux	SIGNATURE OF OBANER	
NAME (Please print or type)	nlug	NAME Please print or type)	
NAME (Please print or type) Daniel W. Gorbet	uluf		
NAME (Please print or type)	DATE April 11, 2003	NAME (Please print or type) Richard L. Jones	ATE 4/25/03

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that is will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

> **Plant Variety Protection Office** Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

18a. Give:

ITEM

(1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;

(2) the details of subsequent stages of selection and multiplication;

- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;

(2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and

- (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.

(Please provide a statement as to the limitation and sequence of generations that may be certified.)

As noted (Breeder, Foundation, Registered, Certified (one year each).

22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

May 2002 (Foundation)

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

Three utility patents effect this variety: 1) No. 5,922,390; 2) No. 6,063,984;

3) No. 6,121,472

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. valid OMB control number for this information collection is 1995, an agency may not conduct or sponsor, and a person is not required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, collider beliefs, parental status, or projected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information. (Braille, large print, audiotape, etc.) should contact USDA STARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

ST-470 (02-10-2003) designed by the Plant Variety Protection Office with Word 2000. Replaces former versions of ST-470, which are a

16a. Exhibit A - Origin and Breeding History of Variety - Hull

Hull (UF98326) came from a three-way cross made in the greenhouse at Marianna, Florida in 1989. The cross was made to incorporate the "high oleic" oil chemistry in material to select for good pod/seed yields, grades, medium-late maturity, resistance to multiple disease (tomato spotted wilt virus, late leafspot/C. personatum, and white mold (S. rolfsii), and improved oil chemistry. A pedigree selection program was followed in the F_1 - F_6 under unsprayed (no leafspot control) conditions. Hull originates from a high oleic seed (HOl) from an F_1 plant from which individual seed were analyzed for oil chemistry. HOl was a "high oleic" seed ($80\pm\%$ 18:1). Single plant selections were made under unsprayed conditions in F_2 - F_6 . Hull was first yield tested at Marianna in 1996 and in Marianna and Gainesville in 1997-2001.

Pedigree = $89 \times OL28-HO1-7-4-1-2-\underline{b3}-B$ [F₁ (Southern Runner x F435-HO) x UF81206]

The female parent was an F₁ plant from a cross of Southern Runner with F435-HO. The F435-HO is a breeding line from the University of Florida breeding program with "high oleic" oil (80±%, 18:1) chemistry of it seed. This line came from an outcross/mutation selection from a Florispan derivative with high oleic chemistry. F435-HO was first reported in a 1987 article in *Peanut Science* (Norden, et al.). F435-HO has been widely used in peanut breeding programs around the world. SunOleic 95R was the first commercial peanut cultivar released with this trait and breeding background. The male parent was UF81206, which is a late maturity University of Florida breeding line that has very good resistance to late leafspot, tomato spotted wilt virus, and white mold. Southern Runner and UF81206 both have PI203396 as a parent. Hull is a jumbo-runner market-type peanut with prostrate growth habit, classified as *Arachis hypogaea* spp. *hypogaea* var. *hypogaea*. Hull has somewhat less vine growth, shorter center stem height and lighter green foliage than C-99R. Seed (testa) of Hull are tan in color and similar in size to those of C-99R.

Hull was first put into yield tests at Marianna in 1996 followed by Gainesville in 1997. Hull has been uniform and stable for all yield tests and seed increases during this period (1996-2004) of evaluation. It has consistently shown to have the high oleic seed trait, multiple disease resistance, and the phenotype described. No variants have been observed.

16.b.c. Exhibit B - Novelty Statement

'Hull' is most similar to 'C-99R'; however, 'Hull' has higher oleic acid (80 vs 57 %) and lower linoleic acid (3 vs 23 %) content in the seed oil than 'C-99R'.

Exhibit C - Objective Description of Variety

Hull is a runner market-type peanut (A. hypogaea L.) with prostrate (runner) growth habit. The foliage is medium to somewhat lighter green than C-99R, especially near maturity. Seed of Hull are similar in size and shape to those of C-99R with a tan testa. The 100-seed weight of Hull is about 70 grams. Pod yields of Hull have been very similar to C-99R in sprayed and unsprayed tests. Hull has resistance to tomato spotted wilt virus, late leafspot (C. personatum), and white mold (S. rolfsii), similar to C-99R, with less vine growth and seed that have "high oleic" oil chemistry. Hull has about 80% oleic (18:1) and 2% linoleic (18:2) fatty acid composition in the seed, with near 50% oil. The oil chemistry is similar to SunOleic 97R.

Additional details to characterize Hull are given in the tables referenced in Exhibit D. Hull has performed very similar to C-99R in many Florida tests.

References:

- 1) Gorbet, Daniel W. 2003. Hull A new multiple disease resistant high oleic peanut variety. UF Agric. Expt. Stn. NFREC Res. Rpt. 03-6. 7 p.
- 2) Gorbet, D. W., and D. A. Knauft. 2000. Registration of 'SunOleic 97R' Peanut. Crop Sci. 40:1190-1191.
- 3) Gorbet, D. W., and F. M. Shokes. 2002. Registration of 'C-99R' Peanut. Crop Sci. 42:2207.

Table 6. Oil chemistry by year and location in Florida tests (1999-2002).

Year/location	·			
variety	Palmitic	Oleic	Linoleic	Oil
1999 Gainesville		% -		
Hull	5.3	81.0	1.7	 .
C-99R	9.8	58.0	22.0	
Fla. MDR 98	4.9	68.1	12.0	
DP-1	7.0	71.2	11.2	
SunOleic 97R	5.3	79.7	3.2	.
1999 Marianna	·			
Hull	5.4	80.5	2.1	49.3
C-99R	9.3	56.3	23.7	51.8
Fla. MDR 98	8.1	62.2	17.9	50.8
DP-1	8.7	61.1	19.4	51.9
SunOleic 97R	6.2	80.3	2.8	50.2
2000 Gainesville				y
Hull	5.8	79.6	2.2	49.1
C-99R	9.7	53.3	24.7	52.1
Fla. MDR 98	8.0	61.5	17.7	53.8
DP-1	8.8	59.0	20.2	50.6
SunOleic 97R	6.1	80.4	2.7	51.8
2000 Marianna				
Hull	5.4	78.3	2.7	48.8
C-99R	8.4	58.7	20.0	52.1
Fla. MDR 98	7.1	66.1	13.3	52.8
DP-1	9.2	57.1	21.0	51.9
SunOleic 97R	5.8	81.0	2.6	50.2
		-		

Table 6 Continued

2001 Gainesville				
Hull	5.9	80.6	3.5	52.0
C-99R	10.0	56.0	23.9	49.6
Fla. MDR 98	8.4	60.6	19.7	53.1
DP-1	9.2	59.4	21.6	52.0
SunOleic 97R	5.9	82.5	3.8	50.9
2001 Marianna			·	
Hull	5.7	80.6	4.2	
C-99R	9.1	57.7	23.5	
DP-1	9.9	53.4	26.9	
SunOleic 97R	5.9	81.3	4.5	·
2002 Gainesville			. •	·
Hull	6.1	80.4	2.1	 ,
C-99R	9.7	57.8	22.6	
DP-1	8.7	62.8	18.6	
SunOleic 97R	6.1	80.8	4.0	_

Table 7. Commercial lab data on chemistry and flavor for Hull (2001).

		Fatty Acids (%)			%		
Entry	16:0	18:1	18:2	Oil	Sugar	Flavor*	
Hull	5.5	79.1	2.9	50.1	3.1	5.2	
C-99R	9.2	53.2	24.8	48.0	3.3	5.0	
Georgia Green	9.7	51.4	28.1	49.3	3.2	4.0	

^{*}Flavor rated on 1-10 (10 = strongest).

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, POULTRY, CRAIN & SEED BIVISION BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY PEANUT (Arachis hypogaea)

NAME OF APPLICANT[3]	VARIETY NAME OR TEMPORARY DESIGNATION
Florida Agricultural Experiment Station ADDRESS (Street and No. of R.F.D. No., City, State, and ZIP Code)	UF98326/Hull
Office of Dean for Research	FOR OFFICIAL USE ONLY
1022 McCarty Hall, University of Florida	200300207
P. O. Box 110200, Gainesville, FL 32611-0200	22 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (e-s. 0 8 9 or 0 9) when number is either 99 or less o	e boxes below. r 9 or less.
1. BOTANICAL TYPE:	
1 Flowering on the Main Stem: 1 = ABSENT 2 = PRESENT	
1 = ALTERNATE - Pairs of vegetative & reproductive branches (Virginal Pattern: 2 = SEQUENTIAL - Continuous reproductive branches (Valencia-Spanners)	
2. PLANT:	A A O D E D A TE (Stare)
1 = PROSTRATE (Florunner) 2 = DECUMBENT (NC-5) Branching: 3 = SEMI-ERECT (Florispan) 4 = ERECT (Starr)	= SPARSE (Valencia) 2 = MODERATE (Starr) 3 = PROFUSE (Florunner)
3. MATURITY:	· ·
Region: 1*VIRGINIA, NORTH CAROLINA 2 = S.E. UNITED STATES 3=S.	W.UNITED STATES 4-OTHER
1 4 5 NUMBER OF DAYS TO MATURITY	
NO. OF DAYS EARLIER THAN	2 = FLORUNNER 3 = FLORIGIANT A 61R 5 = NC - 2 7 = SOUTHEASTERN RUNNER 56-15
1 0 NO. OF DAYS LATER THAN	
4. LEAVES:	
2 COLOR AT 60 DAYS: (Nickerson Color Designation): 1 = LIGHT GREEN (10Gy 3 = DARK GREEN (5G 4/1	
5 7 MM. LEAFLET LENGTH (Basal leaflet of the youngest fully opened leaf)	
2 . 5 LEAFLET LENGTH/WIDTH RATIO	
5. POD: (Average for 20 pods at maturity)	
2 9 MM. LENGTH 1 6 MM. E	DIAMETER
5 3 0 5 KG./HA. POD YIELD	- -
% LESS THAN	2 = FLORUNNER 3 = FLORIGIANT IA 61R 5 = NC - 2
6 = NC-5	7 = SOUTHEASTERN RUNNER 56-15 (Specify) C-99R
6 % FANCY SIZE: (% riding 13.46 mm., 34/64 inch, spacing set on presizer roller)	8

VARIETY OIL (X) PROTEIN* (X) LINOLEIC ACID RATIO IODINE* NUMBER SHELLING (X) SMK** (X) EXX** HEIGHT (CM) SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 51.0 27.0 35.2 76 78,4 77.6 39 33.0 NAME OF SIMILAR VARIETY C-99R C-99R Sun0leic 97R Sun0leic 97R Sun0leic 97R Sun0leic 95R C-99R Georgia Green * From Sound Mature Kernels ** Sound Mature Kernels + Extra Large Kernels ** Sound Mature Kernels ** Sound Mature Kernels ** Sound Mature Kernels										
2 CONSTRICTION: 1 - SHALLOW OR NONE(Viginis 56R, Argentine) 2 SURFACE: 1 - GLASROUS (Fforumer) 2 SURFACE: 1 - GLASROUS (Fforumer) 2 BEAK: 1 - ABSENT 2 - INCONSPICUOUS 3 - PRONOUNCED 5 SEED (Institute, cured but not seed): 1 - WHITE (Pael) 2 - CREAM 3 - TAN (Starr) 4 - BROWN 5 - PINK (Fforighest) 5 - VARIOATED 1 - WARTE (Pael) 1 - COAT SURFACE: 1 - SMOOTH 2 - INDENTED 1 - UNIFORM COLOR 6 - RED 7 - PURPLE 8 - DARK PURPLE 9 - VARIOATED 1 - SPHERIODAL Starr) 2 - SHORT-BROAD (Fforumer) 3 - ELONGATED SLENDER (Deck Rumer) 6 SHAPE: 4 - CYLINORICAL-TAFERED ENDS 5 - CYLINDRICAL-BLUNT ENDS (NC-2) 5 - OTHER (Specify) C-99R 1 S MM, LENGTH 1 O MM, WIDTH 7 O GRAMS PER 100 SEED (S% Moisture) 7. DISEASE RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Repistant) 2 SOUTHERN STEM ROT 1 EARLY LEAF SPOT 0 VIRUS X 2 SOUTHERN STEM ROT 1 EARLY LEAF SPOT 0 WORAIC 0 POD ROT COMPLEX 1 NEMATODE (Specify species) 1 LEAF HOPPER 1 NEMATODE (Specify species) 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 1 NEMATODE (Specify species) 1 NEMA	[•	•							
SURPACE: 1 = GLABROUS (Florunner) 2 = PUBBSCENT (Florighen) 2 = SEAK: 1 - ASSENT 2 = INCONSPICUOUS 3 = PRONOUNCED 6. SEED (Mature, cured but not apped): 6. SEED (Mature, cured but not apped): 6. SEED (Mature, cured but not apped): 7 = PURPLE 8 - CARK PURPLE 9 - VARIGATED 10 - OTHER (Specify) 11 COAT SURPACE: 1 - SMOOTH 2 - INCORNTED 1 1 - UNIFORM COLOR 2 - BLEMISHED 10 - OTHER (Specify) 11 COAT SURPACE: 1 - SMOOTH 2 - INCORNTED 1 1 - UNIFORM COLOR 2 - BLEMISHED 11 - SPHERIODAL (Starr) 2 - SHORT-BROAD (Florunner) 3 - ELONGATED-SLENGER (Disk Runner) 12 SHAPE: 4 - CYLINDRICAL-TAFERED ENDS 5 - CYLINDRICAL-BLUNT ENDS (MC-2) 6 - OTHER (Specify) CC-99R 11 6 MM. LENGTH 1 1 0 MM. WIOTH 7 0 GRAMS PER 100 SEED (# Moldiner) 12 SOUTHERN STEM ROT 1 NUST 13 EARLY LEAF SPOT 0 MOSAIC 14 SOUTHERN STEM ROT 0 VIRUS X 25 SOUTHERN LEAF SPOT 0 MOSAIC 2 SOUTHERN LEAF SPOT 0 MOSAIC 2 SOUTHERN LEAF SPOT 0 MOSAIC 3 SURROWING SUG 1 LESSER CORNSTALK SORER 1 THRIPS 0 BURROWING SUG 1 LESSER CORNSTALK SORER 1 LAPHUD 0 THER (Specify) 2 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK SORER 3 MARIETY (SL) PROTEIN* (MOSAIC) 4 MAIN STEP 1 NEMATODE (SPECIFY) 2 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK SORER 3 MARIETY (SL) PROTEIN* (MICH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 VARIETY (SL) SUBDICAL CORNS (SC) STEEL CHARGETY 1 NOWANCE OF SIMILAR VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 CHARACTER VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 NORDATE A VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 NORDATE A VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 NORDATE A VARIETY WITHCH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 NORDATE A VARIETY WITHCH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 NORDATE A VARIETY WITHCH MOST CLOSELY RESEMBLES THAT SUBMITTED: 1 NORDATE A VARIETY WITHCH MOST CLOSELY RESEMBLES THAT SU	2 NUMBER	OF SEEDS PER	POD: 1=1 2=	2 3=3	4 = 3 -	4 5 = 2-3-4				
EARLY LEAF SPOT Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistance: (0 = Not Tested, 1 = Succeptible, 2 = Resistant) Disease resistant res	2 CONSTRI	CTION: 1 = SHA	LLOW OR NONE(Virg	inia 56R, Arg	entine)	2 = MEDIU	M (Virginia 61R) 3 = D	EEP (Stari)
6. SEED (Mature, cured but not apped): 1 SEED (Mature, cured but not apped): 3 COAT COLOR: 6 - RED 7 - PURPLE 8 - DARK PURPLE 9 - VARIGATED 10 - OTHER (Specify) 1 COAT SURFACE: 1 - SMOOTH 2 - INDENTED 1 1 - UNIFORM COLOR 2 - BLEMISHED 1 - SPHERIODAL (Starr) 2 - SHORT-BROAD (Florumer) 3 - ELONGATED-SLENDER (Dirk Runner) 6 SHAPE: 4 - CYLINDRICAL-TAFERED ENDS 5 - CYLINDRICAL-BLUNT ENDS (NC-2) 6 - OTHER (Specify) C-99R 1 6 MM. LENGTH 1 0 MM. WIDTH 7 0 GRAMS PER 100 SEED (S% Moisture) 7. DISEASE RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) SCALE 1 ; 4 - TCS STATE 1 EARLY LEAF SPOT 0 WIRUS X 2 SOUTHERN STEM ROT 1 RUST 1 EARLY LEAF SPOT 0 MOSAIC 2 OTHER (Specify) Tomato Spotted Wilt Virus 8. INSECT RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) 1 THAIPS 0 BURROWING BUG 1 NEMATOGE (Specify species) 1 LESSER CORNSTALK BORER 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 1 NEMATOGE (Specify species) 2 OTHER (Specify) Tomato Spotted Wilt Virus 3 SUSMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: 1 VARIETY 0 CH. PROTEIN* LINGLE NUMBER (N) SMK** ELK** MEBERTY (CM) 2 SUSMITTED 49,1 26,2 36,5 75 77,5 76,8 39 29,9 3 3.0 OMMARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: 1 VARIETY 0 CH. PROTEIN* LINGLE NUMBER (N) SMC	1 SURFACE	: 1 - GL	ABROUS (Florunner)	2 = PU	BESCENT	(Florispan)	. •			
3 COAT COLOR: 6 - RED 7 - PURPLE 8 - DARK PURPLE 9 - VARIGATED 10 - OTHER Specify - PURPLE 8 - DARK PURPLE 9 - VARIGATED 11 - SPHERIODAL Gram 2 - SHORT-BROAD (Florumer) 3 - ELONGATED-SLENDER (Dick Runner) 11 - SPHERIODAL Gram 2 - SHORT-BROAD (Florumer) 3 - ELONGATED-SLENDER (Dick Runner) 6 SHAPE: 4 - CYLINDRICAL-TAPERED ENDS 5 - CYLINDRICAL-BLUNT ENDS (NC-2) 6 - OTHER (Specify) C-99R 11 6 MM. LENGTH 1 0 MM. WIDTH 7 0 GRAMS PER 100 SEED (#% Molitume) 7. DISEASE RESISTANCE; (0 - Not Tested, 1 - Succeptible, 2 - Resistant) SCAC (-4) 4 - YCS (STAC) 2 SOUTHERN STEM ROT 1 RUST 1 EARLY LEAF SPOT 0 WIRUS X 2 SOUTHERN LEAF SPOT 0 MOSAIC 2 OTHER (Specify) Tomato Spotted Wilt Virus 8. INSECT RESISTANCE; (0 - Not Tested, 1 - Succeptible, 2 - Resistant) 1 THRIPS 0 BURROWING BUG 0 LEAF HOPPER 1 NEMATODE (Specify species) 1 LESSER CORNSTALK BORER 0 APHID 0 THER (Specify) 2 COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY (Nt. PROTEIN ACD TABLE) NUMBER (M. MIMBER M. M. MIMBER M. M. M. M. STEP HEIGHT (SM. M. M. M. M. STEP HEIGHT (SM. M. M. M. M. STEP HEIGHT (SM. M. M. M. M. STEP HEIGHT (SM. M. M. M. M. M. STEP HEIGHT (SM. M. M	2 BEAK:	1 = ABS	SENT 2=	INCONSPICU	ous	3 = PRONO	UNCED			
3 COAT COLOR: 6 - RED 7 - PURPLE 8 - DARK PURPLE 9 - VARIGATED 10 - OTHER (Specify) 10 - OTHER (Specify) 1 - UNIFORM COLOR 2 - BLEMISHED 1 - UNIFORM COLOR 2 - BLEMISHED 1 - SPHERIODAL (Starr) 2 - SHORT-BROAD (Florumer) 3 - ELONGATED SLENDER (Dick Runner) SHAPE: 4 - CYLINDRICAL-TAPERED ENDS 5 - CYLINDRICAL-BLUNT ENDS (NC-2) 6 - OTHER (Specify) C-99R 1 6 MM. LENGTH	6. SEED (Mature, cu	red but not aged):				 		·	
3 COAT COLOR: 6 - RED 10 - OTHER (Specify) 1 - OTHER (Specify) 1 - OTHER (Specify) 1 - OTHER (Specify) 2 - SHORT-BROAD (Florumer) 3 - 6 LONGATED SLEMISHED 1 - UNIFORM COLOR 2 - BLEMISHED 1 - UNIFORM COLOR 2 - BLEMISHED 1 - SHORT-BROAD (Florumer) 3 - 6 LONGATED SLENGER (Dick Rumer) 6 - OTHER (Specify) C - 99R 1 6 MM. LENGTH 1 0 MM. WIDTH 7 0 GRAMS PER 100 SEED (M% Moisture) 7. DISEASE RESISTANCE: (O - Not Tested, 1 - Susceptible, 2 - Resistant) SCALE 4 4 4 7 7 7 7 7 7 7		• • •	WUITE (Donal)	00544	T //			a art. t.t.	41	•
COAT SURFACE: 1 - SMOOTH 2 - INDENTED 1 1 - UNIFORM COLOR 2 - BLEMISHED 1 - SPHERIODAL (Starr) 2 - SHORT-BROAD (Florumer) 3 - ELONGATED-SLENDER (Dick Runner) 3 - ELONGATED-SLENDER (Dick Runner) 6 SHAPE: 4 - CYLINDRICAL-TAPERED ENDS 5 - CYLINDRICAL-BLUNT ENDS (NC-2) 6 - OTHER (Specify) C-99R 1 6 MM. LENGTH 1 0 MM. WIDTH 7 0 GRAMS PER 100 SEED (#% Moisture) 7. DISEASE RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) SCALE 14 4 7 7 7 7 7 7 7 7	3 COAT	COLOR: 6 = 8	RED 7 = 1	PURPLE E		· ·			t)	
SHAPE: 4 - CYLINDRICAL-TAPERED ENDS S - CYLINDRICAL-BUNT ENDS (NC-2) S - OTHER (Specify) C-99R 1 6 MM. LENGTH 1 0 MM. WIDTH 7 0 GRAMS PER 100 SEED (8% Moisture) 7. DISEASE RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) SCOLE 14; 4 - YES SCOLE 15 Moisture 8. SOUTHERN STEM ROT 1 RUST 1 EARLY LEAF SPOT 0 MOSAIC 2 SOUTHERN LEAF SPOT 0 MOSAIC 2 OTHER (Specify) Tomato Spotted Wilt Virus 8. INSECT RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) 1 THRIPS 0 BURROWING BUG 1 LEAF HOPPER 1 NEMATODE (Specify species) 9. COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY 0 IL PROTEIN* LINDLE'C NOMBER (K) SMK** (K) HEIGH (CM) SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 50.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R SUNO 1 C 10 C	1 COAT SUF				1	1 = UNIFORM	COLOR 2	= BLEMIS	HED	
SHAPE: 4 - CYLINDRICAL-TAPERED ENDS S - CYLINDRICAL-BUNT ENDS (NC-2) S - OTHER (Specify) C-99R 1 6 MM. LENGTH 1 0 MM. WIDTH 7 0 GRAMS PER 100 SEED (8% Moisture) 7. DISEASE RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) SCOLE 14; 4 - YES SCOLE 15 Moisture 8. SOUTHERN STEM ROT 1 RUST 1 EARLY LEAF SPOT 0 MOSAIC 2 SOUTHERN LEAF SPOT 0 MOSAIC 2 OTHER (Specify) Tomato Spotted Wilt Virus 8. INSECT RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) 1 THRIPS 0 BURROWING BUG 1 LEAF HOPPER 1 NEMATODE (Specify species) 9. COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY 0 IL PROTEIN* LINDLE'C NOMBER (K) SMK** (K) HEIGH (CM) SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 50.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R SUNO 1 C 10 C		= SPHERIODAL	_ (Starr) 2 = SHC	ORT-BROAD	(Florumne	rl 3=510	NGATED-SI EN	DE <i>B (Divie</i>	Dunner)	
1 6 MM. LENGTH 1 0 MM. WIDTH 7 0 GRAMS PER 100 SEED 18% Moisture) 7. DISEASE RESISTANCE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 1 RUST 1 RUST 1 RUST 2 SOUTHERN STEM ROT 1 POD ROT COMPLEX 2 SOUTHERN LEAF SPOT 0 MOSAIC 2 OTHER (Specify) Tomato Spotted Wilt Virus 8. INSECT RESISTANCE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 1 THRIPS 0 BURROWING BUG 1 LEAF HOPPER 1 NEMATODE (Specify species) 2 OTHER (Specify) 3 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 3 COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY 0 SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 51.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR CAPIETY SUBMITTED: SUBMITTED 49.1 26.2 36.5 75 8 SURDIES SU	1 61									C-99R
7. DISEASE RESISTANCE: (0 - Not Tested, 1 - Susceptible, 2 - Resistant) SCALE 4 - Y CSISTANT 1	·		· · · · · · · · · · · · · · · · · · ·	5 676111	DITICAL	SCORT ERDS DVC	<i>3–2)</i> 6–3	JIHEN WP		
SOUTHERN STEM ROT 1 RUST 1 RUST 2 SOUTHERN LEAF SPOT 0 VIRUS X 2 SOUTHERN LEAF SPOT 0 MOSAIC 2 OTHER (Specify) Tomato Spotted Wilt Virus	1 6 MM. LE	истн []	L 0 MM, WIDT	н 7	0	GRAMS PER 1	00 SEED (8% M	oisture) ′		
BUST BARLY LEAF SPOT DIAM VIRUS X	7. DISEASE RESIST	ANCE: (O = Not	t Tested, 1 = Susceptible	, 2 = Resistan	n) 5c	ale 14.	4= rps	ista.	<u>.</u> +	
2 SOUTHERN LEAF SPOT 0 POD ROT COMPLEX E. INSECT RESISTANCE: (0 - Not Texted, 1 - Susceptible, 2 - Resistant) 1 THRIPS 0 BURROWING BUG 1 LEAF HOPPER 1 NEMATODE (Specify species) 2 OTHER (Specify) 3 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 3 COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY OIL* PROTEIN* LINDLEIC* ACID RATIO NUMBER SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 51.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R SunOleic 97R SunOleic 97R SunOleic 97R SunOleic 97R SunOleic 97R SUNOLEIC* CHARACTER VARIETY CHARACTER C-99R C-9	71 71							13 100	V 1	
Depart Complex B. INSECT RESISTANCE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 1 Thrips 0 BURROWING BUG 1 NEMATODE (Specify species) 0 SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER 0 APHID 3. COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY OIL* PROTEIN* LINGUELC.* LINGUELC.* LINGUELC.* ACID RATIO NUMBER SHELLING SMK.** (K) CM) SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 50.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R SUNO1eic 97R SUNO1eic 97R SUNO1eic 97R From Sound Mature Kernels From So	1 EARLY LE	AF SPOT			0	rinus x	د در			
B. INSECT RESISTANCE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 1 THRIPS 0 BURROWING BUG 1 NEMATODE (Specify species) 1 LESSER CORNSTALK BORER 0 APHID 3. COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY 0 C COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY 0 C COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY 0 C COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY 0 C COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETY 0 C COLOR 1 C SUBMITTED 0 DEIC: 1 C COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETY 0 C COLOR 1 C SUBMITTED 0 DEIC: 1 C C C C C C C C C C C C C C C C C C	2 SOUTHERN	LEAF SPOT			0 ~	IOSAIC				
Description of Submitted Variety with one or more similar varieties: Variety Oil Protein (S) Submitted 49.1 26.2 36.5 75 77.5 76.8 39 29.9 Similar 51.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R Sunoleic 97R Sunoleic 95R Sunoleic 95R C-99R Green From Sound Mature Kernels **Sound Mature Kernels + Extra Large Kernels OILEGE (S) Submitted Variety Withom ost closely resembles that submitted: Character Variety C-99R Seed Link (S) Character C-99R Seed Link (S) Character Variety C-99R Seed Link (S) Character Variety C-99R Seed Link (S) Character C-99R Seed Link (S) Character Variety C-99R Seed Link (S) C-99R Seed Dormancy C-99R Seed Link (S) C-99R Seed Dormancy C-99R Lick (S) C-99R Seed Dormancy C-99R Lick (S) C-99R Seed Dormancy C-99R Lick (S) C-99R Seed Dormancy C-99R Seed Link (S) C-99R Seed L	0 POD ROT C	COMPLEX			2 0	THER (Specify)	Tomato S	potted	Wilt V	<u> irus</u>
O LEAF HOPPER 1 NEMATODE (Specify species) O SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER O APHID OTHER (Specify) O COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY OIL* (K) PROTEIN* LINGLEIC (DODINE* ACIO RATIO NUMBER (K) NUMBER (K) SHELLING SMK** (K) (K) NUMBER (K) NUMBER T7.5 T6.8 39 29.9 SIMILAR 51.0 27.0 35.2 T6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T8.4 T8.6 T8.4 T8.6 T8.6 T8.7 T8	8. INSECT RESISTA	NCE: (0 = Not T	ested, 1 = Susceptible, 2	2 = Resistant)	<u> </u>	. <u>:</u>	·		······	<u> </u>
O LEAF HOPPER 1 NEMATODE (Specify species) O SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER O APHID OTHER (Specify) O COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY OIL* (K) PROTEIN* LINGLEIC (DODINE* ACIO RATIO NUMBER (K) NUMBER (K) SHELLING SMK** (K) (K) NUMBER (K) NUMBER T7.5 T6.8 39 29.9 SIMILAR 51.0 27.0 35.2 T6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T7.6 T8.4 T8.4 T8.6 T8.4 T8.6 T8.6 T8.7 T8	THRIPS		•	-						
O SOUTHERN CORN ROOTWORM 1 LESSER CORNSTALK BORER OTHER (Specify) 3. COMPARISON OF SUBMITTED VARIETY WITH ONE OR MORE SIMILAR VARIETIES: VARIETY OIL PROTEIN LINOLEIC OIDINE SHELLING (X) SMK** (X) HEIGHT (X	\					y kirkin Harriy	Nisasa.			
O APHID OTHER (Specify) OTHER (Specify	U CEAF HOT	CN	•	•	<u> </u>	EMATODE (Spec	ify species)			
ACID RATIO WARIETY OIL® PROTEIN® LINOLEIC FACID RATIO SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR SIMILAR SIMILAR OPPR C-99R SunOleic 97R SunOleic 97R SunOleic 97R SunOleic 97R SunOleic 97R SunOleic 97R C-99R SunOleic 97R CHARACTER WARIETY WARIETY CHARACTER VARIETY CHARACTER VARIETY CHARACTER VARIETY C-99R SEED LING VIGOR C-99R SEED ORMANCY C-99R SEED ORMANCY C-99R SEED ORMANCY C-99R SEED SIZE C-99R LEAF COLOR C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification — Such as: Relative disease reactions may be compared with standard varieties) Hull has good Fesistance to tomato spotted wilt virus and white mold (S. rolfsii) with	0 SOUTHERN	CORN ROOTW	ORM	•		ESSER CORNST	ALK BORER	•		
VARIETY OIL* PROTEIN* (X) PROTEI	0 APHID				$\overline{\square}$ \circ	THER (Specify)	· · · · · · · · · · · · · · · · · · ·			·
VARIETY (%) ACID RATIO NUMBER (%) SMK** (%) (%) HEIGHT (CM) SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 50.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R SunOleic 97R SunOleic 97R SunOleic 97R SunOleic 97R C-99R Green * From Sound Mature Kernels ** Sound Mature Kernels + Extra Large Kernels 10. INDICATE A VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER VARIETY CHARACTER VARIETY COO COLOR C-99R SEEDLING VIGOR C-99R SEED DORMANCY C-99R HULL THICKNESS C-99R SEED DORMANCY C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good Fesistance to tomato spotted wilt virus and white mold (S. rolfsii) with	9. COMPARISON OF	SUBMITTED VA	ARIETY WITH ONE OF	R MORE SIM	LAR VAI	RIETIES:	·			
SUBMITTED 49.1 26.2 36.5 75 77.5 76.8 39 29.9 SIMILAR 5D.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R SunOleic 97R SunOleic 97R SunOleic 97R SunOleic 97R C-99R Green *From Sound Mature Kernels ** Sound Mature Kernels + Extra Large Kernels 10. INDICATE A VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER VARIETY CHARACTER VARIETY COD COLOR C-99R SEEDLING VIGOR C-99R SEED DORMANCY C-99R HULL THICKNESS C-99R SEED SIZE C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with	VARIETY	OIL.	PROTEIN*	LINO	LEIC		SHELLING (%)			! HEIGHT
SIMILAR 50.0 27.0 35.2 76 78.4 77.6 39 33.0 NAME OF SIMILAR C-99R C-99R Sun0leic 97R Sun0leic 97R Sun0leic 97R Sun0leic 97R Sun0leic 97R C-99R Green *From Sound Mature Kernels ** Sound Mature Kernels + Extra Large Kernels 10. INDICATE A VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER VARIETY CHARACTER VARIETY COO COLOR C-99R SEEDLING VIGOR C-99R SEED OORMANCY C-99R HULL THICKNESS C-99R SEED SIZE C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with	SUBMITTED	49.1	26.2	36.5		75	77.5	76.8	39	
*From Sound Mature Kernels ** Sound Mature Kernels + Extra Large Kernels 10. INDICATE A VARIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER VARIETY CHARACTER VARIETY COD COLOR C-99R SEEDLING VIGOR C-99R GEED DORMANCY C-99R HULL THICKNESS C-99R GEED SIZE C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with	SIMILAR	50.0	27.0	35.2		76	78,4	77.6		33.0
* From Sound Mature Kernels ** Sound Mature Kernels + Extra Large Kernels 10. INDICATE A VÁRIETY WHICH MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER VARIETY CHARACTER VARIETY COO COLOR C-99R SEEDLING VIGOR C-99R SEED DORMANCY C-99R HULL THICKNESS C-99R SEED SIZE C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with	NAME OF SIMILAR VARIETY		C-99R	SunOleic	97R	SumOleic 97R		ŧ		
CHARACTER VARIETY CHARACTER VARIETY COO COLOR C-99R SEEDLING VIGOR C-99R SEED DORMANCY C-99R HULL THICKNESS C-99R SEED SIZE C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with						· .				
COO COLOR C—99R SEED LING VIGOR C—99R HULL THICKNESS C—99R SEED SIZE C—99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification — Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with			·	EMBLES THA	T SUBMI					
HULL THICKNESS C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with		TER							RIETY	
EED SIZE C-99R LEAF COLOR Andru 93 11. COMMENTS (Additional description or clarification – Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with			····						·	
11. COMMENTS (Additional description or clarification - Such as: Relative disease reactions may be compared with standard varieties) Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with									~~	
Hull has good resistance to tomato spotted wilt virus and white mold (S. rolfsii) with		ditional description	C-99R					Andru	93	
and only with high officers.	Hull has go	od resista:	nce to tomato	spotted	wilt v	irus and wh	ite mold	(S. rol	fsii)	with

DRM LPGS-470-29 (8-80) (REVERSE)

16d. Exhibit D - Additional Description of Variety

Hull is a medium-late maturity (145 \pm 3 DAP) runner market-type peanut with excellent pod/seed yield potential, very good resistance to tomato spotted wilt virus (TSWV) and white mold (S. rolfsii), moderate resistance to late leafspot (C. personatum) and CBR, with excellent oil quality (high oleic).

Table 1 gives data on pod yield, grades and disease ratings for Florida tests conducted at Marianna and Gainesville (1997-2001). These data show that Hull has pod yields very similar to C-99R under both sprayed and unsprayed conditions. The dominant leafspot pathogen for tests at Marianna is *Cercosporidum personatum* (late leafspot). The data on sprayed tests, comparing Hull to Georgia Green, show a pod yield and disease resistance advantage for Hull. Grades of Hull indicate a somewhat lower SMK value (compared to C-99R and Georgia Green), with an ELK and 100-seed weight almost the same as C-99R.

Table 2 gives data on TSWV tests grown at Marianna, Florida, and Tifton, Georgia, under high pressure conditions (early April planting, four seed/row foot). Hull shows a clear advantage over the resistant check(Georgia Green) for pod yields and disease resistance in all tests. Disease rating was based on percent of plot showing symptoms.

Table 3 gives data on field paired-row tests, where two rows of a four-row plot were inoculated at 50± DAP with *S. rolfsii* (white mold) and two rows were left uninoculated. Results on pod yields and disease ratings indicate that Hull is very similar to C-99R in resistance to this disease and both are more resistant than Georgia Green and Florunner. The disease in the uninoculated rows was mainly TSWV.

Table 4 gives data from Marianna on leafspot resistance studies, comparing Hull to C-99R and Southern Runner. Main-plot treatments in split-plot studies were either unsprayed or sprayed with four fungicide applications on a 21-day schedule, beginning about 55 DAP. Pod yields and disease ratings indicate that Hull has resistance to leafspot (*C. personatum*) very similar to C-99R and better than Southern Runner.

Table 5 gives data on oil chemistry from Florida samples analyzed in University of Florida labs (1999-2000). The results show the "high oleic" (18:1) oil content of Hull, being similar to SunOleic 97R, with the low linoleic (18:2) content. This chemistry has shown to help shelf-life/stability of the oil and for products made from peanuts with this type oil chemistry, compared to "normal" chemistry (Florunner, Georgia Green, C-99R). Data from University of Florida studies have shown health benefits from this type peanut, lowering LDL cholesterol in human

trials in Gainesville.

Table 6 gives data on oil chemistry for Hull compared to C-99R, DP-1, and SunOleic 97R in Gainesville and Marianna Tests. This further shows the high oleic chemistry by location and year, similar to SunOleic 97R.

Table 7 gives additional data on seed chemistry and flavor from 2001 Marianna crop samples, analyzed at a commercial lab. These data further support the "high oleic" oil chemistry of Hull and indicate that it rates ok for roasted flavor.

Table 8 gives blanching data on Florida samples (Marianna, 1998-2000), comparing Hull to Georgia green and C-99R. Hull shows somewhat less whole-seed blanch than Georgia Green and C-99R with somewhat more splits and partials.

Table 9 gives data on seed size distribution for Hull compared to Georgia Green and C-99R at Marianna (1999-2001). Hull is very similar to C-99R in seed size with a somewhat lower meat content. This agrees with the grading data.

Table 1. Pod yield and grading data for Hull in Florida tests (1997-2001).

Entry	Pod Yield (lbs./A)	% Meat	% TSMK¹	% ELK²	100-seed wt. (g)	<u>Dis</u>	ease ³
Sprayed (31) ⁴				· · · · · · · · · · · · · · · · · · ·			
Hull	4737	77.5 -	76.8	39.9	71.9	2.8	3.4
C-99R	4615	. 78.8	78.3	39.6	73.7	2.9	3.4
Unsprayed (21)		•					
Hull	3035	79.9	77.7	36.4	69.2	5.8	2.4
C-99R	3077	81.2	79.4	36.3	69.0	5.6	2.5
Sprayed (21)							
Hull	4644	77.1	75.7	37.5	72.0	3.0	3.3
Georgia Green	3664	79.8	77.8	16.1	57.5	4.2	2.8

¹TSMK = sound mature kernels, seed riding a 16/64th inch slotted screen.

²ELK = extra large kernels, seed riding a 21.5/64th inch slotted screen.

³Disease rated A = 1-10 scale, 1 = no disease (sprayed = TSWV, unsprayed = leafspot), B = 4-1 scale (4 = highly resistant).

⁴Number in parentheses = number of tests.

Table 2. Tomato spotted wilt studies in Florida and Georgia (1998-2000).

		% Disease ¹		Po	d Yield (kg/	ha)
Entry/Year	GA	FL -	Mean	GA	FL	Mean
1998						
Georgia Green	50.8	65.0	57.9	4570	3699	4135
Georgia Runner	81.7	89.6	85.6	2943	2645	2794
Hull	23.3	35.4	29.4	6364	4655	5510
<u>1999</u>						
Georgia Green			50.4	3363	2414	2888
GK 7		· ·	82.7	2081	814	1448
Hull			40.9	4793	4176	4485
2000	•					
Georgia Green	54.6	29.0	41.8	2706	2936	2821
GK 7	62.1	40.0	51.1	2473	1971	2222
Hull	30.8	32.1	31.5	4859	4427	4643

¹Data from Dr. Albert Culbreath, University of Georgia, Tifton.

Table 3. Data on paired-row tests inoculated with S. rolfsii, 1999-2001.

	Yield (kg/ha)	Disease			
Entry	Inoc.	Not	Inoc.1	Inoc. ²	Not^1	Not²
Florunner	2175	3102	7.1	2.0	5.8	. 2.4
Georgia Green	3780	4651	4.3	3.0	3.2	3.3
C-99R	4311	4675	3.0	3.4	2.5	3.7
Hull	<u>4854</u>	<u>5480</u>	<u>3.2</u>	<u>3.4</u>	<u>2.5</u>	<u>3.7</u>
LSD(.05) =	.51	.5		0.3	·	

Disease rated 1-10 (1 = no disease).
Disease rated 1-4 (4 = resistant).

Table 4. Data on leafspot resistance studies at Marianna (1999-2002).

	Pod Yield	%	100-seed	Dis	ease ¹
Entry/Year	(lbs/A)	TSMK	wt. (g)	(1-10)	(1-4)
<u>Unsprayed</u>				•	
Hull	3638	78.1	70.6	5.0	2.6
C-99R	3437	79.9	68.5	5.3	2.6
Southern Runner	2820	78.1	52.9	5.2	2.6
Sprayed ²					
Hull	4875	78.5	70.5	2.8	3.4
C-99R	4689	80.5	72.5	3.3	3.3
Southern Runner	3989	79.4	56.3	3.5	3.2

¹Disease rated 1-10 (1 = no disease); 1-4 (4 = resistant).

Table 5. Oil chemistry data from Florida samples conducted in University of Florida labs (1999-2000)*.

Entry	Oleic (18:1)	Linoleic (18:2)	Oil
		%	
Hull	79.7	2.0	48.8
C-99R	58.8	20.5	52.0
Georgia Green	54.8	25.1	51.3
Florunner	56.0	24.1	49.7
SunOleic 97R	80.7	2.5	49.0

^{*}Data based on no less than 10 fatty acid samples and four samples for oil.

²Sprayed treatment = four fungicide sprays for leafspot control on 21 day schedule, beginning about 55 days.

Table 8. Blanching data on Hull from Marianna samples (1999-2001).

Entry	Splits	Whole	Not	Partial
		%		
Hull	9.7	75.8	4.3	7.5
Georgia Green	7.3	83.7	3.3	3.7
C-99R	4.8	85.0	2.0	5.5

Table 9. Hull seed size distribution data (1998-2000).

	Percent riding screen size						
Entry	21/64"	18/64"	16/64"	14/64"	SS^1	OK²	Meat
·	a a a.	.,		%			
Hull	41.5	19.8	4.2	1.8	7.3	2.0	76.6
C-99R	45.9	20.0	3.0	1.1	6.8	1.7	78.5
Georgia Green	29.8	37.5	5.4	1.7	2.7	1.7	78.8

¹SS = sound splits. ²OK = other kernels (pass thru 14/64).

16.e. Ownership Statement

Hull originates from a 1989 cross made in the greenhouse at the Marianna by D. W. Gorbet. All selections were made at Marianna (D. W. Gorbet) under unsprayed (leafspot) field conditions after the initial analyzing of the F_2 seed (from F_1 plant), for oil chemistry. Hull originates from the HOl (high oleic) F_2 seed. Seed from three F_6 plants were bulked to begin yield testing at Marianna in 1996. Yield tests were continued at Marianna and Gainesville in 1997-2001. UF98326 was approved for release by the University of Florida Agricultural Experiment Station (FAES) in 2002 as a new high oleic, late maturity, multiple disease resistant peanut cultivar, named Hull.

Florida Foundation Seed Producers, Inc. (FFSP) has been designated and authorized to produce breeder and foundation seed of Hull for commercial distribution. Only companies with approved contracts with FFSP are authorized to produce and sell seed of Hull.

Hull was developed by FAES scientist (breeder). By agreement between the breeder and FAES, this invention belongs to FAES and all rights, access, and use of this invention shall be in accordance to FAES policy.

Also, Hull is impacted by three University of Florida utility patents on the "high oleic" trait (Patents # 5,922,030; 6,121,472; and 6,063,984). These three patents impact the commercialization of all high oleic peanuts and as such, arrangements must be make with the University of Florida Foundation for marketing any such peanut varieties.

Public reporting burden for this collection of information is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter.

Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791.

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.